	Faxed prescriptions will only	be accepted from a Gastrointes		ts must bring an original prescription to the pharmacy, and cannot fax these referral forms to Senderra. Prescriber: NPI:							
		Enrollment A-H	Form	Supervising Physician:					NPI:		
SENDERRA Physician C 855-460-79			Offices Call:	Address:					Tax ID:		
Specialty Pharmacy			28	Phone: Fax:							
3712 E. Plano Parkway, Ste. 200 Fax: 888-777-			7-5645	Contact:							
Plano, TX 75074											
	is to be sent & received via fa	x		PATIENT IN	IFORMAT						
Name:			F 🗆 Trans M 🗆 Trans F 🗆 Other				SS#:				
Street: City:				State:				ZIP:	ZIP:		
Phone: Alt. Phone:			□ English □ Spanish □ Other:				Wt.:	Wt.: Ht.:			
PRESCRIPTION											
Has the patient received a loading dose/starter kit? Yes Start Date:// DNo SHIP TO: Patient's Home Doctor's Office Other:											
Drug	П	Directions & Quantity Refills									
Cimzia® □ 200 mg Pre-filled Syringe □ 200 mg Vial			☐ INITIAL: Inject 400 mg (two 200 mg injections) SQ on day 0, 14, and 28 (Quantity: 6) ☐ MAINTENANCE: Inject 400 mg (two 200 mg injections) SQ every 4 weeks (Quantity: 2)								
Dupixent®	300 mg Pre-filled Syringe		Inject 300 mg SQ every week (Quantity: 4) ***WEIGHT REQUIRED*** ***Intended for ages 12 and older with weight ≥ 40 kg/88 lbs***								
			□ INITIAL: Infuse 300 mg IV over 30 minutes at Day 0, 14, and 42 (Quantity: 3)								
Entyvio	Entyvio® 300 mg Vial			MAINTENANCE: Infuse 300 mg IV over 30 minutes every 8 weeks (Quantity: 1)							
	□ 80 mg/0.8 mL Crohn's/UC Starter Kit □ 40 mg/0.4 mL Pen □ 40 mg/0.4 mL Pre-filled Syringe		ADULT: INITIAL: Inject 160 mg SQ on day 1, 80 mg on day 15, then 40 mg every other week starting day 29 (Quantity: 3) MAINTENANCE: Inject 40 mg SQ every other week (Quantity: 2)								
Humira® Citrate Free	☐ 40 mg/0.4 mL Pre-filled Syringe & 20 mg/0.2 mL Pre-filled Syringe		PEDIATRIC: ***WEIGHT REQUIRED*** INITIAL: Inject 80 mg (two 40 mg injections) SQ on day 1, 40 mg (two 20 mg injections) on day 15, then 20 mg every other week starting on day 29 (Quantity: 4)								
	20 mg/0.2 mL Pre-filled Syringe		MAINTENANCE: Inject 20 mg SQ every other week (Quantity: 2) ##Intended for weight 17 kg/37 lbs to <40 kd/88 lbs***								
	□ 80 mg/0.8 mL Crohn's Starter Kit		□ INITIAL: Inject 160 mg SQ on day 1, 80 mg on day 15, then 40 mg every other week starting day 29 (Quantity: 3)								
	40 mg/0.4 mL Pen 40 mg/0.4 mL Pre-filled Syringe		MAINTENANCE: Inject 40 mg SQ every other week (Quantity: 2) ***Intended for weight ≥40 kg/88 lbs***								
			PEDIATRIC: ***WEIGHT REQUIRED***								
	40 mg/0.4 mL Pen 40 mg/0.4 mL Pre-filled Syringe		INITIAL: Inject 80 mg SQ on day 1, 40 mg on day 8, 40 mg on day 15 (Quantity: 4) ***Intended for weight 20 kg/44 lbs								
			MAINTENANCE: Inject 40 mg SQ every other week starting on day 29 (Quantity: 2) to <40 kg/88 lbs***								
	20 mg/0.2 mL Pre-filled Syringe		MAINTENANCE: Inject 20 mg SQ every week starting on day 29 (Quantity: 4)								
	80 mg/0.8 mL Pediatric UC Starter Kit 80 mg/0.8 mL Pen		□ INITIAL: Inject 160 mg SQ on day 1, 80 mg on day 8, 80 mg on day 15 (Quantity: 4)								
	40 mg/0.4 mL Pen		MAINTENANCE: Inject 80 mg SQ every other week starting on day 29 (Quantity: 2) MAINTENANCE: Inject 40 mg SQ every week starting on day 29 (Quantity: 4) MAINTENANCE: Inject 40 mg SQ every week starting on day 29 (Quantity: 4)								
	40 mg/0.4 mL Pre-fille	ed Syringe	☐ MAINTENA				g on day 29 (Quantity	: 4)	_	<u> </u>	
MEDICAL INFORMATION ***PLEASE FAX COPY OF PRESCRIPTION/MEDICAL CARD, FRONT AND BACK, AS WELL AS ANY CLINICAL NOTES REGARDING THERAPY***											
PREVIOUS THE	Not Tolerated:				Con	traindication:					
Methotrexate)								
Sulfasalazine Pentasa)							
Cimzia											
Humira											
la											
K20.0 Eosinop	philic Esophagitis				K20.	<u> </u>					
		tine, without comp	olications			rohn's diseas	e of large intestine,	without co	omplications		
☐ K50.00 Crohn's disease of small intestine, without complications ☐ K50.80 Crohn's disease of both intestines, without complications ☐ K50.90 Crohn's disease unspecified, without complications											
□ K51.50 Left-sided Ulcerative Colitis, without complications □ K51.80 Other Ulcerative Colitis, without complications											
	tive Colitis unspecified,				Other: _						
Date of Diagnosis:// Allergies: Patient is steroid dependent											
Active TB is ruled out:											
Additional Clinical Information:											
INJECTION TRAINING											
□ Patient has received pen and injection training □ Physician's office to provide injection training □ Senderra to coordinate injection training PRESCRIBER SIGNATURE											
To Prescriber: By signing this form and utilizing our services, you are also authorizing Senderra to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies, and co-pay assistance foundations.											
Prescriber:	pay assistance foundations.						Da	te:			
				CONFIDENCE	A1 1734 544	TIOE					
CONFIDENTIALITY NOTICE IMPORTANT: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, proprietary or exempt from disclosure under applicable law. If you are not the named											
addressee, you sho	uld not disseminate, distribut	te, or copy this fax.	Please notify the s	ender immediatel	y if you hav	e received this d	ocument in error and th	nen destroy	this document immediately.		