Faxed prescriptions will only be accepted from a prescribing practitioner. Patients must bring an original prescription to the pharmacy, and cannot fax these referral forms to Senderra.

		Juvenile Idiopathic Arthritis (JIA)	Prescriber:			NPI:	NPI:		
		Enrollment Form	Supervising Physician:			NPI:	NPI:		
SENDERRA Speciality Pharmacy		Physician Offices Call:	Address:			Tax ID:	Tax ID:		
		855-460-7928	Phone: Fax:						
3712 E. Plano Parkway, Ste. 200 Plano, TX 75074		Fax: 888-777-5645	Contact:	1					
This prescription form is to be sent & received via fax PATIENT INFORMATION									
Name:		□ M □ F □ Trans M □ Trans F □ Other DOB:				, SS#:			
Street:		Cit		////////		ZIP:	ZIP:		
Phone:		Alt. Phone:			Wt.:	.: Ht.:			
	Refill Ship by:		SHIP TO: Patient's Home Doctor's Office Other:						
Drug		/	Direc		Quantity Refills				
Diug		INTRAVENOUS (IV):				rtennis			
Orencia®	250 mg Vial WEIGHT REQUIRED:				***WEIGH	T BASED LINES:***	QS: 3 doses		
						5 kg: 10 mg/kg)			
		MAINTENANCE: Infuse mg via IV every 4 weeks				(75 kg-100 kg: 750 mg) (≥100 kg: 1000 mg)			
		SUBCUTANEOUS (SQ):							
	Pre-filled Syringe	□ Inject 50 mg SQ once	***WEIGHT R	***WEIGHT REQUIRED***					
	□ ClickJect ™	□ Inject 87.5 mg SQ once weekly (25 kg to less than 50 kg) □ Inject 125 mg SQ once weekly (≥50 kg)							
Stelara®		\Box INITIAL: Inject mg (0.75 mg/kg xkg) SQ at weeks 0 & 4				***WEIGHT REQUIRED*** QS			
	□ 45 mg Vial	□ MAINTENANCE: Inject mg (0.75 mg/kg xkg) SQ every 12 weeks							
	45 mg Pre-filled Syringe		□ INITIAL: Inject 45 mg SQ at weeks 0 & 4			***Intended for weight ≥ 60 kg/132 lbs***			
		MAINTENANCE: Inject 45 mg SQ every 12 weeks INITIAL: Inject 90 mg SQ at weeks 0 & 4			***Intended for weight	***Intended for weight > 100 kg/220 lbs with co-existent moderate-to-severe plaque psoriasis*** 2			
	□ 90 mg Pre-filled Syringe	MAINTENANCE: Inject 90 mg SQ every 12 weeks		co-existent moderat					
Xeljanz®	5 mg Tablet	□ Take 5 mg PO twice daily		— ***WEIGHT R	- ***WEIGHT REQUIRED*** 				
		☐ Take 3.2 mg PO twice daily (10 kg to less than 20 kg)							
	1 mg/mL Solution	□ Take 4 mg PO twice daily (20 kg to less than 40 kg)							
		□ Take 5 mg PO twice daily (≥40 kg)							
MEDICAL INFORMATION									
PREVIOUS THERAPIES: Tried & Failed (Duration): Not Tolerated: Contraindication:									
		$\exists_{(})$							
		J()					· · · · · · · · · · · · · · · · · · ·		
		رر ۱ ()							
□ Naproxen/Aleve □									
		_(/					· · · · · · · · · · · · · · · · · · ·		
La Humira						· · · · · · · · · · · · · · · · · · ·			
		-()]()				<u></u>			
Date of Diagnosis: / Allergies:									
M08.00 Unspecified Juvenile Idiopathic Arthritis of Unspecified Site M08.09 Unspecified juvenile rheumatoid arthritis, multiple sites (pcJIA)									
L40.54 Psoriatic juvenile arthropathy (JPsA)									
Active TB is ruled out: 🛛 Yes ☐No Date:// Hep B ruled out/treated: ☐Yes ☐No Date://									
Additional Clinical Information:									
INJECTION TRAINING									
Patient has received pen and injection training Physician's office to provide injection training Senderra to coordinate injection training									
PRESCRIBER SIGNATURE To Prescriber: By signing this form and utilizing our services, you are also authorizing Senderra Rx to serve as your prior authorization designated agent in dealing with medical and prescription insurance									
companies, and co-pay assistance foundations. Prescriber: Date:							mouta		
						//	/		
CONFIDENTIALITY NOTICE IMPORTANT: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, proprietary or exempt from disclosure under applicable law. If you are not the named									
addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately.									