					ts must bring an original prescription to the pharmacy, and cannot fax these referral forms to Senderra.								
In		Miscellaneous Immunology			Prescriber:					NPI:			
			rollment Form		Supervising Physician:					NPI:			
SENDERRA Phy		ъ.	sician Offices Call: -460-7928		Address:					Tax ID:			
Specialty Pharmacy		-			Phone: Fax:								
Richardson, TX 75081				Contact:									
This prescription form is to be sent & received via fax:			888-777-5645		- Contact.								
PATIENT INFORMATION Name: DOB: SS#:													
M D F D Trans M D Trans F D Other													
Street:				City:			State:			ZIP:			
Phone: Alt. Ph			ione:		☐ English ☐ Spanish ☐ O		her:		Wt.:		Ht.:		
PRESCRIPTION													
Has the patient received a loading dose/starter kit? Yes Start Date: / / No SHIP TO: Patient's Home Doctor's Office Other: Drug Directions & Quantity Refil													
	☐ 100 mL NS IV bag					Billottolio	a quantity	<u>'</u>					
Infusion Supplies	250 mL NS IV bag												
Actemra [®]	80 mg Vial		Infuse mg OR 8 mg/kg via IV over 1 hour (Quantity: QS 1 dose)										
	400 mg Vial		Infuse mg OR 12 mg/kg via IV over 1 hour (Quantity: QS 1 dose)										
	☐ 120 mg/5 mL Vial		INTRAVENOUS (IV):										
Benlysta [®]	400 mg/20 mL Vial		□ INITIAL: Infusemg or 10 mg/kg via IV over 1 hour every 2 weeks, for 3 doses (Quantity: QS 3 doses) □ MAINTENANCE: Infusemg OR 10 mg/kg via IV over 1 hour every 4 weeks (Quantity: QS 1 dose)										
		SUBCUTANEOUS (SQ):											
, , , ,	200 mg Autoinjector 200 mg Pre-filled syringe		□ INITIAL: Inject 400 mg SQ (two 200 mg injections) once weekly for four weeks (Quantity: 8) ***Dosing intended for Lupus Nephritis***										
			MAINTENANCE: Inject 200 mg SQ every week (Quantity: 4)										
_				☐ Inject 200 mg SQ every week (Quantity: 4) ☐ Inject 300 mg SQ every 4 weeks (Quantity: 3)									
Nucala®	☐ 100 mg Autoinjector		☐ Inject 300 n	ng SQ e	every 4 v	veeks (Quantity: 3)							
Remicade®													
□ _{Avsola®} □ _{Inflectra®}	☐ 100 mg Vial		INITIAL: In	INITIAL: Infuse mg OR mg/kg via IV at weeks 0, 2, and 6 (Quantity: QS 3 doses) MAINTENANCE: Infuse mg OR mg/kg via IV every weeks thereafter (Quantity: QS 1 dose)									
□ Inflectra®	100 mg viai		□ MAINTENA										
Infliximab													
□Rituxan®	☐ 100 mg/10 mL Vial		☐ Infuse	ma o	n 🗆 D:	ay 1 and Day 15 \square Once a week	for 4 week	ks D Other:					
□Ruxience® □ _{Truxima} ®	500 mg/50 mL Vial		100 mg Vial Quantity: 500 mg Vial Quantity:										
	☐ 50 mg Vial		■ INITIAL: Infuse 2 mg/kg via IV over 30 minutes at weeks 0 and 4 (Quantity: QS 2 doses)										
Simponi Aria®	Weight Required:		■ MAINTENANCE: Infuse 2 mg/kg via IV over 30 minutes every 8 weeks thereafter (Quantity: QS 1 dose)										
Simponi Aria	Height Deguired		INITIAL: Infuse 80 mg/m² via IV over 30 minutes at weeks 0 and 4 (Quantity: QS 2 doses) ***Dosing intended for										
	Height Required:		☐ MAINTENA	NCE: Ir		mg/m² via IV over 30 minutes eve	ery 8 week	s thereafter (Quantity	y: QS 1	1 dose)	JIA***		
PLEA	SE FAX COPY OF PRE	ESCRIF	TION/MEDICA	L CAR		CAL INFORMATION <mark>NT AND BACK, AS WELL AS</mark>	ANY CLI	NICAL NOTES RE	GARI	DING TH	IERAPY		
PREVIOUS THERAP		Tried	& Failed (Dura		_,	Not Tolerated:				dication			
☐ Methotrexate ☐ Enbrel)									
Humira)									
<u></u>													
□ C71 Functional disorders of polymorphonuclear neutrophils (CGD) □ D72.119 Hypereosinophilic syndrome (HES), unspecified □ D89.839 Cytokine release syndrome, grade unspecified □ K50.90 Crohn's disease, unspecified, without complications													
L10.0 Pemphigus Vulgaris													
L40.0 Psoriasis Vulgaris L40.0 Psoriasis Vulgaris M05.9 Rheumatoid Arthritis with Rheumatoid Factor, unspecified L40.50 Arthropathic Psoriasis, unspecified (Psoriatic Arthritis)													
	Arthritis with Rheumatoid Juvenile Idiopathic Arthri					M06.9 Rheumatoid Arthritis, un M30.1 Eosinophilic granulomat		olvangiitis (EGPA)					
M31.30 Granulomat	osis with polyangiitis (We	gener's)				M31.7 Microscopic polyangiitis							
	ipus Erythematosus, orga	n or sys	tem involvement	unspe	cified F	M32.14 Glomerular disease in	systemic Iu	ipus erythematosus	(Lupus	Nephritis	s)		
M45.9 Ankylosing Spondylitis, unspecified Q78.2 Osteopetrosis													
Date of Diagnosis:	1 1			Alle	ergies:_								
Active TB is ruled out:	□ _{Yes} □ _I	No	Date:	//		Hep B ruled out/treated:	□ _{Ye}	s D _{No Date}	e:				
Additional Clinical Inf	ormation:					TOTION TO THE STATE OF							
INJECTION TRAINING ☐ Patient has received pen and injection training ☐ Physician's office to provide injection training ☐ Senderra to coordinate injection training													
PRESCRIBER SIGNATURE REQUIREDSTAMPED SIGNATURE NOT ALLOWED													
To Prescriber: By signing this form and utilizing our services, you are also authorizing Senderra to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies, and co-pay assistance foundations.													
PRODUCT SUBSTITUTION PERMITTED DISPENSE AS WRITTEN													
X			Date:			X DENTIALITY NOTICE				Date:			
						erial that is confidential, proprietary mediately if you have received this d						named	