Faxed prescriptions will only be accepted from a prescriber. Patients must bring an original prescription to the pharmacy, and cannot fax these referral forms to Senderra.  Adalimumab Biosimilar   Prescriber:   NPI:										
	3		Adalimumab Biosimilar Abrilada		r:	NPI:				
			Amjevita Cyltezo		ng Physician:	NPI:				
SENDERRA  Specialty Pharmacy		Enrollm	Enrollment Form			Tax ID:				
						Fax:				
Physicia 3712 E. Plano Parkway, Ste. 200			n Offices Call: 7928	Phone:						
Plano, TX 75074										
This prescription form is to be sent & received via fax  Fax: 888-777-5645  PATIENT INFORMATION										
Name:			□ M □ F □ Trans M □ Trans F □ Other □ DOB:					1	SS#:	
Street:			City: State:				· ·	ZIP:		
Phone:	A	It. Phone:	☐ English ☐ Spanish ☐ Other:				Wt.:	Ht.:		
PRESCRIPTION										
Has the patient received a loading dose/starter kit? Yes Start Date: / / No Ship to: Patient's Home Doctor's Office Other:  Drug Directions & Quantity										
Abrilada <sup>™</sup>	□ 40 mg/0.8 mL Pen □ 40 mg/0.8 mL Pre-filled Syringe □ 20 mg/0.4 mL Pre-filled Syringe		INITIAL/LOADING DOSES:  ***WEIGHT REQUIRED  ***  PSORIASIS/UVEITIS: Inject 80 mg on SQ day 1, 40 mg on day 8, then 40 mg every other week (Quantity: 4)  CROHN'S/UC/HS: Inject 160 mg on SQ on day 1, then 80 mg on day 15 (Quantity: 6)  ***Intended for ped CD patients ≥ 40kg (88 lbs)***  PEDIATRIC CROHN'S: Inject 80 mg on SQ on day 1, 40 mg on day 15, then 20 mg every other week starting on day 29 (Quantity: QS)  MAINTENANCE DOSES:  Inject 40 mg SQ every other week (Quantity: 2)  Inject 40 mg SQ weekly (Quantity: 4)  Inject 80 mg SQ every other week (Quantity: 4)  Inject 20 mg SQ every other week (Quantity: 2)  ***Intended for JIA patients 15 kg (33 lbs) to <30 kg (66 lbs) OR Crohn's disease patients 17kg (37 lbs) to loss than 40 kg (88 lbs)***							
Amjevita <sup>®</sup>	B0 mg/0.8 mL Sureclick® Autoinjector 40 mg/0.4 mL Sureclick® Autoinjector Pre-filled Syringe 40 mg/0.4 mL 20 mg/0.4 mL 20 mg/0.2 mL 10mg/0.2 mL		INITIAL/LOADING DOSES:  "***WEIGHT REQUIRED****  □ PSORIASIS/UVEITIS: Inject 80 mg on SQ day 1, 40 mg on day 8, then 40 mg every other week (Quantity: QS 28 Days)  □ CROHN'S/UC/HS: Inject 160 mg on SQ on day 1, then 80 mg on day 15 (Quantity: QS 28 Days)  □ PEDIATRIC CROHN'S: Inject 80 mg on SQ on day 1, then 40 mg on day 15 (Quantity: QS 28 Days)  ■ MAINTENANCE DOSES: □ Inject 40 mg SQ every other week (Quantity: 2) □ Inject 40 mg SQ weekly (Quantity: 4) □ Inject 80 mg SQ every other week (Quantity: 2) □ Inject 10 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2) □ Inject 20 mg SQ every other week (Quantity: 2)							
Cyltezo®	□ 40 mg/0.8 mL Pen □ 40 mg/0.4 mL Pen □ Pre-filled Syringe □ 40 mg/0.8 mL □ 40 mg/0.4 mL □ 20 mg/0.4 mL □ 10mg/0.2mL		INITIAL/LOADING  PSORIASIS/UV  CROHN'S/UC/H  PEDIATRIC CR  MAINTENANCE D  Inject 40 mg SQ  Inject 40 mg SQ  Inject 80 mg SQ  Inject 10 mg SQ	DOSES: EITIS: Inject IS: Inject 160 OHN'S: Inject OSES: every other weekly (Qua every other every other	***WE 80 mg on SQ day 1, 0 mg on SQ on day 1 ct 80 mg on SQ on d week (Quantity: 2) antity: 4) week (Quantity: 4)	40 mg of the state	O mg on day 15 (G en 40 mg on day 1	Quantity: 6) 5 (Quantity: 3) g (22 lbs) to <15 kg (33 lbs	***Intended for ped CD patients ≥ 40kg (88 lbs)***  ***Intended for weight 17 kg (37 lbs) to < 40 kg (88 lbs)***	
MEDICAL INFORMATION  ***PLEASE FAX COPY OF PRESCRIPTION/MEDICAL CARD, FRONT AND BACK, AS WELL AS ANY CLINICAL NOTES REGARDING THERAPY***										
PREVIOUS T			I (Duration):	AND, PROP	Not Toler		AU ANT CLINIC		ntraindication:	
☐ Methotrexa	ate		) )		_ _ _					
L40.0 Psoriasis Vulgaris (Plaque Psoriasis)  M06.9 Rheumatoid Arthritis, unspecified  M08.09 Unspecified juvenile RA, multiple sites (pcJIA)  M45.9 Ankylosing Spondylitis, unspecified  M45.9 Ankylosing Spondylitis, unspecified										
Date of Diagnosis:    //										
Active TB is ruled out: Uyes Uno Date: / / Hep B ruled out/treated: Uyes Uno Date: / / Additional Clinical Information:										
				IN IE	ECTION TRAINING					
Patient has received pen and injection training Physician's office to provide injection training Senderra to coordinate injection training										
To Prescriber	By signing this form and utilizing or	ur services	you are also authorizin		CRIBER SIGNATUR		on designated ager	nt in dealing with me	edical and prescription insurance of	ompanies
To Prescriber: By signing this form and utilizing our services, you are also authorizing Senderra to serve as your prior authorization designated agent in dealing with medical and prescription insurance companiand co-pay assistance foundations.  Prescriber:  Date:										
				CONFI	DENTIALITY NOTIC	E				
IMPORTANT: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, proprietary or exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately.										