



**Gastrointestinal Enrollment Form A-H**  
**Physician Offices Call: 855-460-7928**  
**Fax: 888-777-5645**

3712 E. Plano Parkway, Ste. 200  
 Plano, TX 75074  
*This prescription form is to be sent & received via fax*

<b>Prescriber:</b>	<b>NPI:</b>
<b>Supervising Physician:</b>	<b>NPI:</b>
<b>Address:</b>	<b>Tax ID:</b>
<b>Phone:</b>	<b>Fax:</b>
<b>Contact:</b>	

**PATIENT INFORMATION**

<b>Name:</b>	<input type="checkbox"/> M <input type="checkbox"/> F <input type="checkbox"/> Trans M <input type="checkbox"/> Trans F <input type="checkbox"/> Other	<b>DOB:</b> / /	<b>SS#:</b> - -
<b>Street:</b>	<b>City:</b>	<b>State:</b>	<b>ZIP:</b>
<b>Phone:</b>	<b>Alt. Phone:</b>	<input type="checkbox"/> English <input type="checkbox"/> Spanish <input type="checkbox"/> Other: _____	<b>Wt.:</b> <b>Ht.:</b>

**PRESCRIPTION**

Has the patient received a loading dose/starter kit?  Yes Start Date: \_\_\_/\_\_\_/\_\_\_  No **SHIP TO:**  Patient's Home  Doctor's Office  Other: \_\_\_\_\_

Drug	Directions & Quantity	Refills
<b>Cimzia®</b> <input type="checkbox"/> 200 mg Pre-filled Syringe <input type="checkbox"/> 200 mg Vial	<input type="checkbox"/> <b>INITIAL:</b> Inject 400 mg (two 200 mg injections) SQ on day 0, 14, and 28 (Quantity: 6) <input type="checkbox"/> <b>MAINTENANCE:</b> Inject 400 mg (two 200 mg injections) SQ every 4 weeks (Quantity: 2)	
<b>Dupixent®</b> <input type="checkbox"/> 200 mg Pre-filled Syringe <input type="checkbox"/> 200 mg Pen* <input type="checkbox"/> 300 mg Pre-filled Syringe <input type="checkbox"/> 300 mg Pen*	<b>***WEIGHT REQUIRED***</b> _____ <b>*** Dupixent pen indicated for ages 2 and older*</b> <input type="checkbox"/> Inject 200 mg SQ every <b>other</b> week (Quantity: 2) <b>***Intended for ages 1 and older with weight 15kg/33 lbs to &lt; 30 kg/66 lbs***</b> <input type="checkbox"/> Inject 300 mg SQ every <b>other</b> week (Quantity: 2) <b>***Intended for ages 1 and older with weight 30kg/66 lbs to &lt; 40 kg/88 lbs***</b> <input type="checkbox"/> Inject 300 mg SQ every week (Quantity: 4) <b>***Intended for ages 1 and older with weight ≥ 40 kg/88 lbs***</b>	
<b>Entyvio®</b> <input type="checkbox"/> 300 mg Vial	<input type="checkbox"/> <b>INITIAL:</b> Infuse 300 mg IV over 30 minutes at Day 0, 14, and 42 (Quantity: 3) <input type="checkbox"/> <b>MAINTENANCE:</b> Infuse 300 mg IV over 30 minutes every 8 weeks (Quantity: 1)	
<b>Humira® Citrate Free</b> <input type="checkbox"/> 80 mg/0.8 mL Crohn's/UC Starter Kit <input type="checkbox"/> 40 mg/0.4 mL Pen <input type="checkbox"/> 40 mg/0.4 mL Pre-filled Syringe <input type="checkbox"/> 40 mg/0.4 mL Pre-filled Syringe & 20 mg/0.2 mL Pre-filled Syringe <input type="checkbox"/> 20 mg/0.2 mL Pre-filled Syringe <input type="checkbox"/> 80 mg/0.8 mL Crohn's Starter Kit <input type="checkbox"/> 40 mg/0.4 mL Pen <input type="checkbox"/> 40 mg/0.4 mL Pre-filled Syringe <input type="checkbox"/> 40 mg/0.4 mL Pen <input type="checkbox"/> 40 mg/0.4 mL Pre-filled Syringe <input type="checkbox"/> 20 mg/0.2 mL Pre-filled Syringe <input type="checkbox"/> 80 mg/0.8 mL Pediatric UC Starter Kit <input type="checkbox"/> 80 mg/0.8 mL Pen <input type="checkbox"/> 40 mg/0.4 mL Pen <input type="checkbox"/> 40 mg/0.4 mL Pre-filled Syringe	<b>ADULT:</b> <input type="checkbox"/> <b>INITIAL:</b> Inject 160 mg SQ on day 1, 80 mg on day 15, then 40 mg every <b>other</b> week starting day 29 (Quantity: 3) <input type="checkbox"/> <b>MAINTENANCE:</b> Inject 40 mg SQ every <b>other</b> week (Quantity: 2)	
	<b>PEDIATRIC: ***WEIGHT REQUIRED***</b> _____ <input type="checkbox"/> <b>INITIAL:</b> Inject 80 mg (two 40 mg injections) SQ on day 1, 40 mg (two 20 mg injections) on day 15, then 20 mg every <b>other</b> week starting on day 29 (Quantity: 4) <input type="checkbox"/> <b>MAINTENANCE:</b> Inject 20 mg SQ every <b>other</b> week (Quantity: 2) <b>***Intended for weight 17 kg/37 lbs to &lt;40 kg/88 lbs***</b>	
	<input type="checkbox"/> <b>INITIAL:</b> Inject 160 mg SQ on day 1, 80 mg on day 15, then 40 mg every <b>other</b> week starting day 29 (Quantity: 3) <input type="checkbox"/> <b>MAINTENANCE:</b> Inject 40 mg SQ every <b>other</b> week (Quantity: 2) <b>***Intended for weight ≥40 kg/88 lbs***</b>	
	<b>PEDIATRIC: ***WEIGHT REQUIRED***</b> _____ <input type="checkbox"/> <b>INITIAL:</b> Inject 80 mg SQ on day 1, 40 mg on day 8, 40 mg on day 15 (Quantity: 4) <b>***Intended for weight 20 kg/44 lbs to &lt;40 kg/88 lbs***</b> <input type="checkbox"/> <b>MAINTENANCE:</b> Inject 40 mg SQ every <b>other</b> week starting on day 29 (Quantity: 2)	
	<input type="checkbox"/> <b>MAINTENANCE:</b> Inject 20 mg SQ every week starting on day 29 (Quantity: 4)	
	<input type="checkbox"/> <b>INITIAL:</b> Inject 160 mg SQ on day 1, 80 mg on day 8, 80 mg on day 15 (Quantity: 4) <input type="checkbox"/> <b>MAINTENANCE:</b> Inject 80 mg SQ every <b>other</b> week starting on day 29 (Quantity: 2) <b>***Intended for weight ≥40 kg/88 lbs***</b> <input type="checkbox"/> <b>MAINTENANCE:</b> Inject 40 mg SQ every week starting on day 29 (Quantity: 4)	

**MEDICAL INFORMATION**

**\*\*\*PLEASE FAX COPY OF PRESCRIPTION/MEDICAL CARD, FRONT AND BACK, AS WELL AS ANY CLINICAL NOTES REGARDING THERAPY\*\*\***

<b>PREVIOUS THERAPIES:</b>	<b>Tried &amp; Failed (Duration):</b>	<b>Not Tolerated:</b>	<b>Contraindication:</b>
<input type="checkbox"/> Methotrexate	<input type="checkbox"/> (_____)	<input type="checkbox"/>	_____
<input type="checkbox"/> Pentasa	<input type="checkbox"/> (_____)	<input type="checkbox"/>	_____
<input type="checkbox"/> Cimzia	<input type="checkbox"/> (_____)	<input type="checkbox"/>	_____
<input type="checkbox"/> Humira	<input type="checkbox"/> (_____)	<input type="checkbox"/>	_____
<input type="checkbox"/> _____	<input type="checkbox"/> (_____)	<input type="checkbox"/>	_____
<input type="checkbox"/> _____	<input type="checkbox"/> (_____)	<input type="checkbox"/>	_____

<input type="checkbox"/> K20.0 Eosinophilic Esophagitis	<input type="checkbox"/> K20. _____
<input type="checkbox"/> K50.00 Crohn's disease of small intestine, without complications	<input type="checkbox"/> K50.10 Crohn's disease of large intestine, without complications
<input type="checkbox"/> K50.80 Crohn's disease of both intestines, without complications	<input type="checkbox"/> K50.90 Crohn's disease unspecified, without complications
<input type="checkbox"/> K51.50 Left-sided Ulcerative Colitis, without complications	<input type="checkbox"/> K51.80 Other Ulcerative Colitis, without complications
<input type="checkbox"/> K51.90 Ulcerative Colitis unspecified, without complications	<input type="checkbox"/> Other: _____

**Date of Diagnosis:** \_\_\_/\_\_\_/\_\_\_ **Allergies:** \_\_\_\_\_  Patient is steroid dependent  
 Active TB is ruled out:  Yes  No **Date:** \_\_\_/\_\_\_/\_\_\_ Hep B ruled out/treated:  Yes  No **Date:** \_\_\_/\_\_\_/\_\_\_

**Additional Clinical Information:**

**INJECTION TRAINING**

Patient has received pen and injection training  Physician's office to provide injection training  Senderra to coordinate injection training

**PRESCRIBER SIGNATURE**

**To Prescriber:** By signing this form and utilizing our services, you are also authorizing Senderra to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies, and co-pay assistance foundations.

**Prescriber:** \_\_\_\_\_ **Date:** \_\_\_/\_\_\_/\_\_\_

**CONFIDENTIALITY NOTICE**

**IMPORTANT:** This fax is intended to be delivered only to the named addressee. It contains material that is confidential, proprietary or exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately.